datasheet NS Reagents



Anti-HDAC6 Antibody

Alternative Names: CPBHM, HD6, JM21, PPP1R90, Histone deacetylase 6, Tubulin-lysine deacetylase HDAC6

Catalogue Number: AB19-10114-100ug

Size: 100 µg

Background Information

Histone deacetylase 6 (HDAC6) is a member of the HDAC family of enzymes that remove acetyl groups from lysines of histones thereby acting as gene transcription regulators. HDAC6 is more specifically a member of the class IIB HDAC family and contains two catalytic domains and a C-terminal zinc finger domain that binds free ubiquitin as well as ubiquitinated proteins with high affinity. As a cytoplasmic deacetylase, HDAC6 can deacetylate substrates such as tubulin, heat shock protein 90 (HSP90).

HDAC6 plays an important role in protein degradation, alongside roles in cell morphology, adhesion, migration, and tumor cell invasion/metastasis.

Inhibition of Histone deacetylases (HDACs) has been shown to be neuroprotective in animal models of neurodegenerative diseases and HDAC6 is one possible target, however theoretically HDAC6 should actually contribute to the degradation of protein aggregates seen in neurodegenerative disorders such as Alzheimer's, Parkinson's and Hutington's diseases, So HDAC6 as a target for neurodegenerative disorders remains controversial.

Product Information

Antibody Type: Polyclonal Host: Rabbit

Species Reactivity: Human, Mouse Isotype: lgG

Immunogen: Recombinant human HDAC6

 $100 \mu g$ in $100 \mu l$ PBS with 0.02% sodium azide, 50% glycerol, pH7.3. Format:

Storage Conditions: Store at -20°C. Avoid freeze / thaw cycles.

Applications: IHC IF

IHC 1:50-200. IF 1:50-200.

Additional Information

131kDa (Intended as a general **Subcellular location:** Cytoplasm Nucleus Perikaryon axon dendrite MW:

> guide and does not allow for all isoforms and species variations)

Gene ID 10013 **Uniprot ID:** Q9UBN7

tel: +44(0)1638 551500 | fax: +44(0)1638 590481 | tech@nktscientific.com | newmarketscientific.com