

HDAC6 Rabbit pAb

Catalog No.: A11259 **4 Publications**

Basic Information

Observed MW

160kDa

Calculated MW

114kDa/131kDa

Category

Primary antibody

Applications

WB,IHC,IF/ICC,IP

Cross-Reactivity

Human, Mouse, Rat

Background

Histones play a critical role in transcriptional regulation, cell cycle progression, and developmental events. Histone acetylation/deacetylation alters chromosome structure and affects transcription factor access to DNA. The protein encoded by this gene belongs to class II of the histone deacetylase/acuc/apha family. It contains an internal duplication of two catalytic domains which appear to function independently of each other. This protein possesses histone deacetylase activity and represses transcription.

Recommended Dilutions

WB	1:500 - 1:2000
IHC	1:50 - 1:200
IF/ICC	1:50 - 1:200
IP	1:20 - 1:50

Immunogen Information

Gene ID	Swiss Prot
10013	Q9UBN7

Immunogen

Recombinant fusion protein containing a sequence corresponding to amino acids 836-1104 of human HDAC6 (NP_006035.2).

Synonyms

HDAC6;CPBHM;HD6;JM21;PPP1R90

Contact

 | www.abclonal.com

Product Information


Source	Isotype	Purification
Rabbit	IgG	Affinity purification

Storage


Store at -20°C. Avoid freeze / thaw cycles.

Buffer: PBS with 0.02% sodium azide,50% glycerol,pH7.3.


Validation Data




Western blot analysis of extracts of Rat liver, using HDAC6 antibody (A11259) at 1:500 dilution.
Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution.
Lysates/proteins: 25ug per lane.
Blocking buffer: 3% nonfat dry milk in TBST.
Detection: ECL Basic Kit (RM00020).
Exposure time: 1s.



Immunoprecipitation analysis of 100ug extracts of HepG2 cells using 3ug HDAC6 antibody (A11259).
Western blot was performed from the immunoprecipitate using HDAC6 antibody (A11259) at a dilution of 1:1000.



Immunohistochemistry of paraffin-embedded human liver using HDAC6 Rabbit pAb (A11259) at dilution of 1:400 (40x lens). Perform high pressure antigen retrieval with 10 mM citrate buffer pH 6.0 before commencing with IHC staining protocol.



Immunofluorescence analysis of HeLa cells using HDAC6 antibody (A11259) at dilution of 1:100 (40x lens). Blue: DAPI for nuclear staining.