

Affinity Biosciences website:www.affbiotech.com order:order@affbiotech.com

SIRT7 Ab

References(4) Images(10)

Cat.#: DF6161 Concn.: ~1mg/ml Mol.Wt.: 45kDa Size: 100ul,200ul,50ul Source: Rabbit Clonality: Polyclonal

Application: WB 1:500-1:2000, IHC 1:50-1:200, ELISA(peptide) 1:20000-1:40000

*The optimal dilutions should be determined by the end user.

Reactivity: Human, Mouse, Rat, Monkey

Purification: The antiserum was purified by peptide affinity chromatography using

SulfoLinkTM Coupling Resin (Thermo Fisher Scientific).

Immunogen: A synthesized peptide derived from human SIRT7, corresponding to a

region within N-terminal amino acids.

Uniprot: Q9NRC8

Description: This gene encodes a member of the sirtuin family of proteins, homologs to

the yeast Sir2 protein. Members of the sirtuin family are characterized by a sirtuin core domain and grouped into four classes. The functions of human sirtuins have not yet been determined; however, yeast sirtuin proteins are known to regulate epigenetic gene silencing and suppress recombination of rDNA. Studies suggest that the human sirtuins may function as intracellular regulatory proteins with mono-ADP-ribosyltransferase activity. The protein encoded by this gene is included in class IV of the sirtuin family. [provided

by RefSeq, Jul 2008]

Storage: Rabbit IgG in phosphate buffered saline, pH 7.4, 150mM NaCl, 0.02%

sodium azide and 50% glycerol. Store at -20 °C. Stable for 12 months from

date of receipt.



Western blot analysis of extracts from VERO cells, using SIRT7 Ab. The lane

on the left was treated with blocking peptide.



Affinity Biosciences

website:www.affbiotech.com order:order@affbiotech.com



DF6161 at 1/100 staining Rat brain tissue by IHC-P. The sample was formaldehyde fixed and a heat mediated antigen retrieval step in citrate buffer was performed. The sample was then blocked and incubated with the primary Ab at 4°C overnight. An HRP conjugated anti-Rabbit Ab was used as the secondary Ab.

<code>IMPORTANT:</code> For western blot, incubate membrane with diluted primary Ab in 5% w/v milk , 1% TBS, 0.1% Tween@20 at 4°C with gentle shaking, overnight.

For Research Use Only. Not for use in diagnostic and therapeutic procedures. Not for resale without express authorization.