


## PIBF1 Ab

Affinity Biosciences website:www.affbiotech.com order:order@affbiotech.com

Images(2)

Cat.#: DF7171 Size: 100ul,200ul,50ul	Concn.: ~1mg/ml Source: Rabbit	Mol.Wt.: 90kDa Clonality: Polyclonal
Application:	WB 1:500-1:1000, IHC 1:50-1:100, IF/ICC 1:100-1:500, ELISA(peptide) 1:20000-1:40000 *The optimal dilutions should be determined by the end user. Human,Mouse,Rat	
Reactivity:		
Purification:	The antiserum was purified by peptide affinity chromatography using SulfoLink <sup>™</sup> Coupling Resin (Thermo Fisher Scientific).	
Immunogen:	A synthesized peptide derived from hur region within C-terminal amino acids.	nan PIBF1, corresponding to a
Uniprot:	Q8WXW3	
Description:	Progesterone-induced-blocking factor 1 is a protein that in humans is encoded by the PIBF1 gene. It has been shown to localize to the centrosome and has also been named CEP90. Mediator of progesterone that by acting on the phospholipase A2 enzyme interferes with arachidonic acid metabolism, induces a Th2 biased immune response, and by controlling NK activity exerts an anti-abortive effect.	
Storage:	Rabbit IgG in phosphate buffered saline , pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol. Store at -20 $^\circ$ C. Stable for 12 months from date of receipt.	



1 2

250 — 150 —

Western blot analysis of extracts from hepg2, using PIBF1 Ab. Lane 1 was treated with the blocking peptide.

DF7171 staining Hela cells by IF/ICC. The samples were fixed with PFA and permeabilized in 0.1% Triton X-100,then blocked in 10% serum for 45 minutes at 25°C. Samples were then incubated with primary Ab(DF7171 1:200) and mouse anti-beta tubulin Ab(T0023 1:200) for 1 hour at 37°C. An AlexaFluor594 conjugated goat anti-rabbit IgG(H+L) Ab(Red) and an AlexaFluor488 conjugated goat anti-mouse IgG(H+L) Ab(Green) were used as the secondary Ab.

The nuclear counter stain is DAPI(blue).



Affinity Biosciences website:www.affbiotech.com order:order@affbiotech.com

IMPORTANT: For western blot, incubate membrane with diluted primary Ab in 5% w/v milk , 1X TBS, 0.1% Tween®20 at 4°C with gentle shaking, overnight.

For Research Use Only. Not for use in diagnostic and therapeutic procedures. Not for resale without express authorization.